

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

03269/100M292-US3

Figure 1. Analysis of Pain Intensity Difference (PID) Scores Over Time






Figure 2. Analysis of Pain Relief (PR) Scores over Time



03269/100M292-US3

Figure 3. Pooled Efficacy Results on Combined Pain Relief and Pain Intensity Difference (PRID)



03269/100M292-US3

Average (\pm SD) Total Ibuprofen Plasma Concentrations
After Administration of Both Treatments




Figure 4

03269/100M292-US3

Average (\pm SD) Oxycodone Plasma Concentration (ng/mL)
After Administration of Both Treatments






Figure 5



Effects of increasing concentrations of ibuprofen on the permeability of oxycodone across Caco-2 cell monolayers.

* - Significance level: $p < 0.05$, when compared with the permeability value in the absence of ibuprofen.


Figure 6



Effects of increasing concentrations of ibuprofen on the amount of oxycodone transported across Caco-2 cell monolayers after the initial 20 minute-transport period of time.

* - Significance level: $p < 0.05$, when compared with the value in the absence of ibuprofen.

Figure 7



Effects of increasing concentrations of oxycodone on the permeability of ibuprofen across Caco-2 cell monolayers.

Figure 8





Figure 9




◆: One tablet of 5 mg oxycodone hydrochloride/400 mg ibuprofen

■: Two tablets of Nuprin® (200 mg ibuprofen per tablet)

▲: Two tablets of Nuprin® (200 mg ibuprofen per tablet) +
one tablet of Roxicodone™ (5 mg oxycodone hydrochloride).


Figure 10

03269/100M292-US3



- ◆: One tablet of 5 mg oxycodone hydrochloride/400 mg ibuprofen
- : Two tablets of Nuprin[®] (200 mg ibuprofen per tablet)
- ▲: Two tablets of Nuprin[®] (200 mg ibuprofen per tablet) +
one tablet of Roxicodone[™] (5 mg oxycodone hydrochloride).

Figure 11




◆: One tablet of 5 mg oxycodone hydrochloride/400 mg ibuprofen

■: One tablet of Roxicodone™ (5 mg oxycodone hydrochloride)

▲: Two tablets of Nuprin® (200 mg ibuprofen per tablet) +
one tablet of Roxicodone™ (5 mg oxycodone hydrochloride).

Figure 12



◆: One tablet of 5 mg oxycodone hydrochloride/400 mg ibuprofen

■: One tablet of Roxicodone™ (5 mg oxycodone hydrochloride)

▲: Two tablets of Nuprin® (200 mg ibuprofen per tablet) +
one tablet of Roxicodone™ (5 mg oxycodone hydrochloride).

Figure 13